

## Math 18.06 Exam 2 Solutions

**1 (36 pts.)**

(a)  $q_4^* = v - (q_1^T v)q_1 - (q_2^T v)q_2 - (q_3^T v)q_3$   
 $q_4 = \frac{q_4^*}{\|q_4^*\|}$

(b) The nullspace of  $Q$  is just the zero vector ( $Q$  has a pivot in every column). The nullspace of  $Q^T$  has dimension one and consists of all scalar multiples of  $q_4$  (because we know  $q_4$  is orthogonal to  $q_1, q_2$  and  $q_3$ ).

The nullspace of  $Q^T Q = I$  is just the zero vector. The nullspace of  $Q Q^T$  again has dimension one and is all scalar multiples of  $q_4$ .

(c)  $Q^T Q \bar{x} = Q^T b$  is the same as  $\bar{x} = Q^T b$ , so

$$\bar{x} = \begin{bmatrix} q_1^T (q_1 + 2q_2 + 3q_3 + 4q_4) \\ q_2^T (q_1 + 2q_2 + 3q_3 + 4q_4) \\ q_3^T (q_1 + 2q_2 + 3q_3 + 4q_4) \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

The projection  $p = Q \bar{x} = q_1 + 2q_2 + 3q_3$

2 (24 pts.)

(a)  $\begin{bmatrix} 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} x = \begin{bmatrix} 3 \\ 5 \\ K \end{bmatrix}$

$Ax = b$  has an exact solution when  $b$  is in the column space. This happens when  $K = 7$ .

(b)  $\bar{x} = 0$  is the least squares solution when  $b$  is in the nullspace of  $A^T$

For  $\begin{bmatrix} 3 \\ 5 \\ K \end{bmatrix}$  to be in the nullspace of  $A^T$ ,  $K$  would have to be  $-8$  and  $-\frac{21}{4}$ , which is impossible.

- 3 (40 pts.)**
- (a) The determinant will have the cofactor of  $a_{14}$  added to it. In the second part of the question, the determinant will double.
  - (b) We know  $P^2 = P$ , so  $(\det(P))^2 = \det(P)$ , so  $\det(P) = 0$  or  $1$ .
  - (c) Using cofactors by the first row,  $\det(C) = (-b)(-b)(a^2 - b^2) + (-a)(a)(a^2 - b^2) = -(a^2 - b^2)^2$
  - (d) 24 terms using  $a_{11}$  + 24 terms using  $a_{22}$  - 6 terms using both = 42 total