

Your PRINTED name is: SOLUTIONS

Grading

1

2

3

Please circle your recitation:

- 1) M 2 2-131 P. Lee 2-087 2-1193 lee
- 2) M 2 2-132 T. Lawson 4-182 8-6895 tlawson
- 4) T 10 2-132 P.-O. Persson 2-363A 3-4989 persson
- 5) T 11 2-131 P.-O. Persson 2-363A 3-4989 persson
- 6) T 11 2-132 P. Pylyavskyy 2-333 3-7826 pasha
- 7) T 12 2-132 T. Lawson 4-182 8-6895 tlawson
- 8) T 12 2-131 P. Pylyavskyy 2-333 3-7826 pasha
- 9) T 1 2-132 A. Chan 2-588 3-4110 alicec
- 10) T 1 2-131 D. Chebikin 2-333 3-7826 chebikin
- 11) T 2 2-132 A. Chan 2-588 3-4110 alicec
- 12) T 3 2-132 T. Lawson 4-182 8-6895 tlawson

1 (30 pts.) Suppose A is m by n with **linearly dependent columns**. Complete with as much true information as possible:

(a) The rank of A is ...

at most $n - 1$ (and at most m , which is a stronger statement if $m < n - 1$).

(b) The nullspace of A contains ...

at least one non-zero vector. (The dimension of the nullspace is n minus the column rank of A , i.e., at least 1.)

(c) (more words needed) The equation $A^T y = b$ has no solution for some right hand sides b because ...

the rows of the matrix A^T , which are the same as the columns of A , are linearly dependent, so A^T is not full row-rank. Thus the reduced row echelon form of A^T contains a row of all zeroes, so the components of b must satisfy a certain linear relation in order for $A^T y = b$ to have a solution.

2 (40 pts.) Suppose A is this 3 by 4 matrix:

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 6 \end{bmatrix}$$

(a) A specific basis for the column space of A is _____.

(b) For which vectors $b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ does $Ax = b$ have a solution? Give conditions on b_1, b_2, b_3 .

(c) There is no 4 by 3 matrix B for which $AB = I$ (3 by 3). Give a good reason (is this because A is rectangular?).

(d) Find the complete solution to $Ax = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$.

Solution

(a) $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$ is a basis for the column space of A . So is $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$.

(b) Row reducing the augmented matrix $[A \ b]$, we get

$$\left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & b_1 \\ 0 & -1 & -2 & -3 & b_2 - 2b_1 \\ 0 & 0 & 0 & 0 & b_3 - 2b_2 + b_1 \end{array} \right].$$

The linear equations corresponding to the top two rows can be satisfied regardless of the values of b_1 and b_2 , and the bottom row of all zeroes imposes the condition $b_3 - 2b_2 + b_1 = 0$. Hence $Ax = b$ has a solution if and only if $b_3 - 2b_2 + b_1 = 0$.

(c) This is because A is not full row-rank, as shown in part (b). If $AB = I$ then we could solve $Ab_1 = \text{row 1 of } I$, $Ab_2 = \text{row 2 of } I$, $Ab_3 = \text{row 3 of } I$, and every equation $Ax = b$. Actually the solution would be $x = Bb$. But in part (b) we saw that $Ax = b$ has no solution for some b .

(d) We perform the row reduction

$$\left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & 1 \\ 2 & 3 & 4 & 5 & 0 \\ 3 & 4 & 5 & 6 & -1 \end{array} \right] \longrightarrow \left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & 1 \\ 0 & -1 & -2 & -3 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{cccc|c} 1 & 0 & -1 & -2 & -3 \\ 0 & 1 & 2 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Then $x_p = \begin{bmatrix} -3 \\ 2 \\ 0 \\ 0 \end{bmatrix}$ is a particular solution, and $s_3 = \begin{bmatrix} 1 \\ -2 \\ 1 \\ 0 \end{bmatrix}$ and $s_4 = \begin{bmatrix} 2 \\ -3 \\ 0 \\ 1 \end{bmatrix}$ are

special solutions forming a basis of the nullspace of A . Hence the general solution is

$$x = x_p + x_n = x_p + cs_3 + ds_4.$$

- 3 (30 pts.)**
- (a) Find a basis for the vector space of all real 3 by 3 symmetric matrices.
- (b) Suppose A is a square invertible matrix. You permute its rows by a permutation P to get a new matrix B . How do you know that B is also invertible?
- (c) “If 2 matrices have the same shape and the same nullspace, then they have the same column space.” **If this is true**, give a reason why. **If this is not true**, find 2 matrices to show it’s false.

Solution

- (a) The most natural basis is

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

- (b) A being invertible means that A has full rank. Permuting the rows has no effect on the rank, so B has full rank as well, and is thus invertible. (Another argument: the permutation matrix P is invertible, and so $B^{-1} = (PA)^{-1} = A^{-1}P^{-1}$.)

- (c) The statement is false. Example: $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ have the same nullspace (the line spanned by the vector $(1, -1)$), but their column spaces differ (for A , it’s the line spanned by the vector $(1, 1)$, and for B , it’s the line spanned by the vector $(1, 2)$).

Remark (now on the web page)

The real 3 by 3 matrices form a vector space M . The symmetric matrices in M form a subspace S . If you add two symmetric matrices, or multiply by real numbers, the result is still a symmetric matrix. **Problem: Find a basis for S .**

When I asked this question on an exam, I realized that a key point needs to be emphasized: **The basis “vectors” for S must lie in the subspace.** They are 3 by 3 symmetric matrices! Then there are two requirements:

1. The basis vectors must be linearly independent.
2. Their combinations must produce every vector (matrix) in S .

Here is one possible basis (all symmetric) for this example:

$$\begin{array}{ccc} S_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} & S_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} & S_3 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ S_4 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} & S_5 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} & S_6 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \end{array}$$

Since this basis contains 6 vectors, the **dimension of S is 6.**