

**Grading**

1

2

3

4

**Your name is:** \_\_\_\_\_**Please circle your recitation:** \_\_\_\_\_

- 1) M2 2-131 I. Ben-Yaacov 2-101 3-3299 pezz
- 2) M3 2-131 I. Ben-Yaacov 2-101 3-3299 pezz
- 3) M3 2-132 A. Oblomkov 2-092 3-6228 oblomkov
- 4) T11 2-132 A. Oblomkov 2-092 3-6228 oblomkov
- 5) T12 2-132 I. Pak 2-390 3-4390 pak
- 6) T1 2-131 B. Santoro 2-085 2-1192 bsantoro
- 7) T1 2-132 I. Pak 2-390 3-4390 pak
- 8) T2 2-132 B. Santoro 2-085 2-1192 bsantoro
- 9) T2 2-131 J. Santos 2-180 3-4350 jsantos

1 (20 pts.) We are given two vectors  $a$  and  $b$  in  $\mathbb{R}^4$ :

$$a = \begin{bmatrix} 2 \\ 5 \\ 2 \\ 4 \end{bmatrix} \quad b = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}$$

- (a) Find the projection  $p$  of the vector  $b$  onto the line through  $a$ . **Check (!)** that the error  $e = b - p$  is perpendicular to (what?)
- (b) The subspace  $S$  of all vectors in  $\mathbb{R}^4$  that are *perpendicular to this  $a$*  is 3-dimensional. Find the projection  $q$  of  $b$  onto this perpendicular subspace  $S$ . The numerical answer (it doesn't need a big computation!) is  $q = \underline{\hspace{2cm}}$ .



**2 (30 pts.)** Suppose  $q_1, q_2, q_3$  are 3 orthonormal vectors in  $\mathbb{R}^n$ . They go in the columns of an  $n$  by 3 matrix  $Q$ .

(a) What inequality do you know for  $n$ ?

Is there any condition on  $n$  for  $Q^T Q = I$  (3 by 3)?

Is there any condition on  $n$  for  $Q Q^T = I$  ( $n$  by  $n$ )?

(b) Give a nice matrix formula involving  $b$  and  $Q$ , for the projection  $p$  of a vector  $b$  onto the column space of  $Q$ .

**Complete this sentence:**  $p$  is the closest vector \_\_\_\_\_ . . .

(c) Suppose the projection of  $b$  onto that column space is  $p = c_1 q_1 + c_2 q_2 + c_3 q_3$ . Find a formula for  $c_1$  that only involves  $b$  and  $q_1$ . (You could take dot products with  $q_1$ .)



- 3 (20 pts.)** Suppose the 4 by 4 matrix  $M$  has four equal rows all containing  $a, b, c, d$ . We know that  $\det(M) = 0$ . The problem is to find by any method

$$\det(I + M) = \begin{vmatrix} 1+a & b & c & d \\ a & 1+b & c & d \\ a & b & 1+c & d \\ a & b & c & 1+d \end{vmatrix}.$$

**Note** If you can't find  $\det(I + M)$  in general, partial credit for the determinant when  $a = b = c = d = 1$ .

- 4 (30 pts.) We are looking for the parabola  $y = C + Dt + Et^2$  that gives the least squares fit to these four measurements:

$$y_1 = 1 \text{ at } t_1 = -2, \quad y_2 = 1 \text{ at } t_2 = -1, \quad y_3 = 1 \text{ at } t_3 = 1, \quad y_4 = 0 \text{ at } t_4 = 2.$$

- (a) Write down the four equations (not solvable!) for the parabola  $C + Dt + Et^2$  to go through those four points. This is the system  $Ax = b$  to solve by least squares:

$$A \begin{bmatrix} C \\ D \\ E \end{bmatrix} = b.$$

What equations would you solve to find the best  $C, D, E$ ?

- (b) Compute  $A^T A$ . Compute its determinant. Compute its inverse. NOT ASKING FOR  $C, D, E$ .
- (c) The first two columns of  $A$  are already orthogonal. From column 3, subtract its projection onto the plane of the first two columns. That produces what third orthogonal vector  $v$ ? Normalize  $v$  to find the third orthonormal vector  $q_3$  from Gram-Schmidt.

