

Grading

1

2

3

Your name is: SOLUTIONS**Please circle your recitation:**

- 1) M2 2-131 I. Ben-Yaacov 2-101 3-3299 pezz
- 2) M3 2-131 I. Ben-Yaacov 2-101 3-3299 pezz
- 3) M3 2-132 A. Oblomkov 2-092 3-6228 oblomkov
- 4) T11 2-132 A. Oblomkov 2-092 3-6228 oblomkov
- 5) T12 2-132 I. Pak 2-390 3-4390 pak
- 6) T1 2-131 B. Santoro 2-085 2-1192 bsantoro
- 7) T1 2-132 I. Pak 2-390 3-4390 pak
- 8) T2 2-132 B. Santoro 2-085 2-1192 bsantoro
- 9) T2 2-131 J. Santos 2-180 3-4350 jsantos

1 (40 pts.) This question deals with the following symmetric matrix A :

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

One eigenvalue is $\lambda = 1$ with the line of eigenvectors $x = (c, c, 0)$.

- (a) That line is the nullspace of what matrix constructed from A ?
- (b) Find (in any way) the other two eigenvalues of A and two corresponding eigenvectors.
- (c) The diagonalization $A = SAS^{-1}$ has a specially nice form because $A = A^T$. Write all entries in the three matrices in the nice symmetric diagonalization of A .
- (d) Give a reason why e^A is or is not a symmetric positive definite matrix.

Solution:

- (a) The eigenvectors for $\lambda = 1$ make up the nullspace of $A - I$.
- (b) First method: A has trace 2 and determinant -2 . So the two eigenvalues after $\lambda_1 = 1$ will add to 1 and multiply to -2 . Those are $\lambda_2 = 2$ and $\lambda_3 = -1$.

Second method: Compute $\det(A - \lambda I) = -\lambda^3 + 2\lambda^2 + \lambda - 2$ and find the roots 1, 2, -1 : (divide by $\lambda - 1$ to get $\lambda^2 - \lambda - 2 = 0$ for the roots λ_2 and λ_3).

Eigenvectors: $\lambda_2 = 2$ has $x_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$, $\lambda_3 = -1$ has $x_3 = \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}$.

- (c) Every symmetric matrix has the nice form $A = Q\Lambda Q^T$ with orthogonal matrix Q . The columns of Q are orthonormal eigenvectors. (They could be multiplied by -1 .)

$$Q = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{3} & 1/\sqrt{6} \\ 1/\sqrt{2} & -1/\sqrt{3} & -1/\sqrt{6} \\ 0 & 1/\sqrt{3} & -2/\sqrt{6} \end{bmatrix} \quad \text{and} \quad \Lambda = \begin{bmatrix} 1 & & \\ & 2 & \\ & & -1 \end{bmatrix}.$$

- (d) e^A is symmetric and all its eigenvalues e^λ are positive—so e^A is positive definite.

- 2 (30 pts.) (a) Find the *eigenvalues* and *eigenvectors* (depending on c) of

$$A = \begin{bmatrix} .3 & c \\ .7 & 1 - c \end{bmatrix}.$$

For which value of c is the matrix A *not diagonalizable* (so $A = SAS^{-1}$ is impossible)?

- (b) What is the *largest range of values of c* (real number) so that A^n approaches a limiting matrix A^∞ as $n \rightarrow \infty$?
- (c) What is that limit of A^n (still depending on c)? You could work from $A = SAS^{-1}$ to find A^n .

Solution:

- (a) Both columns add to 1. As we know for Markov matrices, $\lambda = 1$ is an eigenvalue. From $\text{trace}(A) = .3 + (1 - c)$ the other eigenvalue is $\lambda = .3 - c$. Check: $\det A = \lambda_1 \lambda_2 = (1)(.3 - c)$ is correct.

The eigenvector for $\lambda = 1$ is in the nullspaces of

$$\begin{aligned} A - I &= \begin{bmatrix} -.7 & c \\ .7 & -c \end{bmatrix} & \text{so } x_1 &= \begin{bmatrix} c \\ .7 \end{bmatrix} \\ A - (.3 - c)I &= \begin{bmatrix} c & c \\ .7 & .7 \end{bmatrix} & \text{so } x_2 &= \begin{bmatrix} 1 \\ -1 \end{bmatrix}. \end{aligned}$$

A is *not diagonalizable* when its eigenvalues are equal: $1 = .3 - c$ or $c = -.7$. (The two eigenvectors above become dependent at $c = -.7$)

(b) $A^n = SAS^{-1} = S \begin{bmatrix} 1 & 0 \\ 0 & (.3 - c)^n \end{bmatrix} S^{-1}$

This approaches a limit if $|.3 - c| < 1$. You could write that out as $-.7 < c < 1.3$ (Small note: at $c = -.7$ the eigenvalues are 1 and 1, at $c = 1.3$ the eigenvalues are 1 and -1 .)

(c) The eigenvectors are in S . As $n \rightarrow \infty$ the smaller eigenvalue λ_2^n goes to zero, leaving

$$\begin{aligned} A^\infty = S \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} S^{-1} &= \begin{bmatrix} c & 1 \\ .7 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ .7 & -c \end{bmatrix} / (c + .7) \\ &= \begin{bmatrix} c \\ .7 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} / (c + .7) = \begin{bmatrix} c & c \\ .7 & .7 \end{bmatrix} / (c + .7) \end{aligned}$$

- 3 (30 pts.) Suppose A (3 by 4) has the Singular Value Decomposition (with real orthogonal matrices U and V)

$$A = U\Sigma V^T = \begin{bmatrix} & & & \\ u_1 & u_2 & u_3 & \\ & & & \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} & & & \\ v_1 & v_2 & v_3 & v_4 \\ & & & \end{bmatrix}^T.$$

- (a) Find the *rank* of A and a *basis* for its column space $C(A)$.
- (b) What are the eigenvalues and eigenvectors of $A^T A$? (You could first multiply A^T times A .)
- (c) What is Av_1 ? You could start with $V^T v_1$ and then multiply by Σ and U to get $U\Sigma V^T v_1$.

Solution:

- (a) Rank = 2 = rank($A^T A$) = # of nonzero singular values. The vectors u_1 and u_2 (very sorry about the typo) are a basis for the column space of A .
- (b) $A^T A = (V\Sigma^T U^T)(U\Sigma V^T) = V\Sigma^T \Sigma V^T$. The eigenvalues of $A^T A$ are 4, 1, 0, 0 in the diagonal matrix $\Sigma^T \Sigma$. The eigenvectors are v_1, v_2, v_3, v_4 in the matrix V .

(c) $V^T v_1 = \begin{bmatrix} v_1^T \\ v_2^T \\ v_3^T \\ v_4^T \end{bmatrix} \begin{bmatrix} \\ \\ v_1 \\ \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$ by orthogonality of the v 's.

Multiply by Σ to get $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$. Then multiply by U to get the final answer $2u_1$.

Thus $Av_1 = 2u_1$, which was a main point of the SVD.