

## 18.06 - Spring 2005 - Problem Set 5

This problem set is due Wednesday (March 16th), at 4 PM, in 2-106. Make sure to PRINT your **name, recitation number and instructor** on your homework!

Please staple your MATLAB solutions as first pages of your homework.

Lecture 15:

- **Read:** book section 4.3.
- **Work:** book section 4.3 (exercises 4, 9, 12, 26 and 27)

Lecture 16:

- **Read:** book section 4.4.
- **Work:** book section 4.4 (exercises 3, 7, 15, 18, 24 and 36).

Lecture 17:

- **Read:** book section 5.1.
- **Work:** book section 5.1 (exercises 3, 12, 15, 28 and 34).

### Challenge Problem with MATLAB

The command `a=ones(n,1)` produces an  $n \times 1$  matrix of 1's.

The command `l=(1:n)'` produces the vector  $(1, 2, \dots, n)$ , transposed to a column by `'`.

The command `s=1.^2` produces the vector  $(1^2, 2^2, \dots, n^2)^T$ , because the dot means "a component at a time."

This problem looks for the line  $y = c + dt$  closest to the parabola  $y = t^2$  on the interval  $t = 0$  to  $t = 1$ .

1. Find the best line by calculus, not MATLAB. Choose  $c$  and  $d$  to minimize

$$E(c, d) := \int_0^1 (c + dt - t^2)^2 dt$$

2. With  $n = 10$ , choose  $C$  and  $D$  to give the line  $y = C + Dt$  that is closest to  $t^2$  at the points  $t = \frac{1}{10}, \frac{2}{10}, \dots, 1$  (in the vector `l/10`). The unsolvable equations  $AX = b$  (use least squares) are

$$\begin{bmatrix} a & l/n \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = s/n^2$$

Find the best  $C$  and  $D$  and the errors  $c - C$  and  $d - D$ .

3. Repeat for  $n = 20$ . (Notice how `l/n` and `s/n^2` end at 1, like the calculus problem.)

Are the differences  $c - C$  and  $d - D$  smaller for  $n = 20$  and by approximately what factor?