

Recitation 6. April 13

Focus: computing determinants, Cramer's rule, diagonalization, eigenvalues and eigenvectors

There are three main ways of computing the determinant of an $n \times n$ matrix A :

- **row echelon form**: row reduce the matrix A , and then:

$$\det A = \pm \text{product of pivots}$$

where the sign is $+$ if you did an even number of row exchanges, and $-$ if you did an odd number of row exchanges.

- **the big formula**:

$$\det A = \sum_{\{\sigma(1), \dots, \sigma(n)\}}^{\text{permutations}} (-1)^{\text{sgn } \sigma} a_{1\sigma(1)} \dots a_{n\sigma(n)}$$

- **cofactor expansion**:

$$\text{along the } i\text{-th row: } \det A = a_{i1}C_{i1} + \dots + a_{in}C_{in}$$

$$\text{along the } i\text{-th column: } \det A = a_{1i}C_{1i} + \dots + a_{ni}C_{ni}$$

where $C_{ij} = (-1)^{i+j}$ times the determinant of the matrix obtained by removing row i and column j from A .

The formulas above also give rise to **cofactor formulas** for inverse matrices:

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} C_{11} & \dots & C_{n1} \\ \vdots & \ddots & \vdots \\ C_{1n} & \dots & C_{nn} \end{bmatrix}$$

The only formula for determinants that you may give without justification is the 2×2 case:

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

Cramer's rule gives a quick formula for the solutions of a system $A\mathbf{v} = \mathbf{b}$ for an $n \times n$ matrix A :

$$\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} \quad \text{where } v_i = \frac{\det B_i}{\det A}$$

and B_i is obtained from A by replacing its i -th column with the vector \mathbf{b} .

To **diagonalize** a square matrix A means to write it as:

$$A = V \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix} V^{-1}$$

Explicitly, the numbers $\lambda_1, \dots, \lambda_n$ are called **eigenvalues** and the columns of V are called **eigenvectors**:

$$V = [\mathbf{v}_1 \mid \dots \mid \mathbf{v}_n]$$

The way you compute these is the following. Eigenvalues are the roots of the characteristic polynomial:

$$p(\lambda) = \det(A - \lambda I)$$

Once you know the eigenvalues, the eigenvectors are computed as bases for nullspaces:

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i \quad \Leftrightarrow \quad \mathbf{v}_i \in N(A - \lambda_i I)$$

1. Compute the determinant of:

$$\begin{bmatrix} 1 & 2 & -1 & 0 \\ 3 & -2 & 0 & 5 \\ -2 & 0 & -2 & 1 \\ 1 & 0 & -1 & 4 \end{bmatrix}$$

by doing a cofactor expansion along its second row.

Solution:

2. Use the cofactor formula to invert the following matrix:

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

Solution:

3. Use Cramer's rule to solve the following system of equations:

$$\begin{cases} x + 3y - z = 0 \\ x + y + 4z = 0 \\ x + z = 1 \end{cases}$$

Solution:

4. Find the eigenvalues and eigenvectors of the following matrix

$$A = \begin{bmatrix} 2 & -1 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

Let $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation given by $\phi(v) = Av$. Can you find a basis v_1, v_2, v_3 of \mathbb{R}^3 with respect to which ϕ is given by a diagonal matrix?

Solution: