18.06 Professor Strang Exam 3 May 6, 1998

Your name is: ______________________________

Grading 1
2
3
4

Please circle your recitation:

1)	M2	2-132	M. Nevins	2-588	3-4110	monica@math
2)	M3	2-131	A. Voronov	2-246	3-3299	voronov@math
3)	T10	2-132	A. Edelman	2-380	3-7770	edelman@math
4)	T12	2-132	A. Edelman	2-380	3-7770	edelman@math
5)	T12	2-131	Z. Spasojevic	2-101	3-4470	zoran@math
6)	T1	2-131	Z. Spasojevic	2-101	3-4770	zoran@math
7)	T2	2-132	Y. Ma	2-333	3-7826	yanyuan@math

1 Find the eigenvalues and eigenvectors of these matrices:

(a) **(10)** Projection $P = \dfrac{aa^T}{a^Ta}$ with $a = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$

(b) **(10)** Rotation $Q = \begin{bmatrix} .6 & -.8 \\ .8 & .6 \end{bmatrix}$

(c) **(8)** Reflection $R = 2P - I$

2 (a) **(10)** Find the eigenvalues λ_1, λ_2, λ_3 (NOT the eigenvectors x_1, x_2, x_3) of this Markov matrix:

$$A = \begin{bmatrix} .6 & .6 & 0 \\ .2 & .2 & .2 \\ .2 & .2 & .8 \end{bmatrix}$$

(b) **(10)** Suppose u_0 is the sum $x_1 + x_2 + x_3$ of the three eigenvectors that you didn't compute. What is $A^n u_0$?

(c) **(4)** As $n \to \infty$ what is the limit of $A^n u_0$?

3 (a) **(2 each)** Suppose M is any invertible matrix. Circle all the properties of a matrix A that remain the same for $M^{-1}AM$:

same rank

same nullspace

same determinant

real eigenvalues

orthonormal eigenvectors

symmetric positive definiteness

(b) **(2 each)** This is a similar question but now Q is an orthonormal matrix. Circle the properties of A that remain the same for $Q^{-1}AQ$:

same column space

A^k approaches zero as k increases

orthonormal eigenvectors

symmetric positive definiteness

projection matrix

4 (a) **(3 each)** Suppose the 5 by 4 matrix A has independent columns. What is the most information you can give about

the eigenvalues of $A^T A$: ____________________

the eigenvectors of $A^T A$: ____________________

the determinant of $A^T A$: ____________________

(b) **(9)** Find the singular value decomposition (SVD) for this matrix:

$$A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 4 & 0 \end{bmatrix} .$$

(c) **(8)** When the input basis is $v_1, \ldots, v_n$ and the output basis is $w_1, \ldots, w_n$ and the matrix of the linear transformation T using these bases is the identity matrix, what is $T(v_1 + v_2)$?