

NEW INDIGO CHROMOPHORES CONTAINING DISULFIDE DONOR GROUPS

Ralf Gerke,^a Lutz Fitjer,^{*a} Peter Müller,^b Isabel Usón,^b Klaus Kowski^c and Paul Rademacher^c

Institut für Organische Chemie der Universität Göttingen,^a Tammannstrasse 2, D-37077 Göttingen, Germany
 Institut für Anorganische Chemie der Universität Göttingen,^b Tammannstrasse 4, D-37077 Göttingen, Germany
 Institut für Organische Chemie der Universität Gesamthochschule Essen,^c Universitätsstrasse 5, D-45141 Essen, Germany

Received 10 September 1999; accepted 11 October 1999

Abstract: Oxidative dimerization of a mixture of the thiolanone **5** and the dithiolanone **4** yields **2c** as chromophore of thioindigo **1c**, and **6** and **7** as chromophores of still unknown indigo dyes. As compared to **2c**, the first ionization potential of the new indigo chromophore **7** is higher, but the calculated HOMO-LUMO splitting is lower. Accordingly, its longest wavelength absorption maximum surpasses that of **2c** distinctly. © 1999 Elsevier Science Ltd. All rights reserved.

Introduction

Predicted by HMO- und PPP-calculations,¹ and confirmed experimentally,²⁻⁴ the 22 π -electron system of the indigo dyes **1a-c** may be reduced to a 10 π -electron system as in **2a-c** without appreciable loss of colour. Thus, **2a-c** represent the basic chromophores of **1a-c** and contain them in the planar arrangement⁵ and the *trans-s-cis-s-cis* configuration characteristic for **1a-c**. As a consequence, their longest wavelength absorption maxima only slightly fall short of those of the corresponding parent compounds **1a-c** (Table 1).





Table 1. Longest wavelength absorption maxima of **1a-c** and **2a-c** in ethanol.

	X	λ_{\max} [nm]		ν_{\max} [cm ⁻¹]	
		1	2	1	2
a	NH	606	480	16 600	20 800
b	Se	559	478	17 890	20 920
c	S	542	453	18 500	22 100

Given the fact that **2a-c** are well tailored to mimic the properties of **1a-c**, it is tempting to devise chromophores of still unknown indigo dyes by just varying the donor and/or acceptor groups. We herein report on the synthesis, structure and spectroscopic properties of two such systems derived from **2c** by incorporating one and two disulfide groups, respectively. As compared to **2c**, the resulting new chromophores **6** and **7** absorb at distinctly longer wavelengths.

Syntheses and Crystal Structures

We obtained **6** (Fp 144°C) and **7** (Fp 191°C) together with **2c**⁴ by oxidative dimerization of a 1:1-mixture of 4,4-dimethyl-thiolan-3-one (**5**)⁶ and 3,3-dimethyl-[1,2]dithiolan-4-one (**4**),⁷ itself obtained by reaction of 1,3-dibromo-3-methyl-butan-2-one (**3**)⁸ with sodium disulfide. Both compounds were separated from each other and from **2c** by column chromatography on silica gel in chloroform and crystallized from ethanol yielding crystals suitable for X-ray crystal structure analyses. These analyses revealed, that both **6** and **7** adopt the *trans-s-cis-s-cis* configuration already established for **2c**.⁴ In the case of **7**, the same conclusion could have been drawn from the IR spectrum. It shows only one (antisymmetrical) carbonyl frequency at 1667 cm⁻¹.



The single-crystal X-ray structures show **6** and **7** to crystallize in the monoclinic space group $P2_1/n$ with half a molecule in the asymmetric unit. The rests of the molecules are generated by the inversion center. In the case of the asymmetric species **6** the molecule is disordered about the inversion center (occupancies 50:50) to fulfil the symmetry of the space group. In both **6** and **7** the five-membered rings adopt a half-chair conformation. These half-chairs are connected such, that the atoms of the central double bond and the adjacent carbon and sulfur atoms take part of one plane, while the peripheral carbon and sulfur atoms, and, to a smaller extent, the oxygen atoms deviate from this plane pairwise in opposite direction. As a result, **6** exhibits a pseudo inversion center, and **7** a real one. Plots of the X-ray crystal structures of **6** and **7** are given in Figure 1.





Figure 1. Crystal structure of **6** (left) and **7** (right) with 50% probability ellipsoids.

UV and PE Spectroscopic Data

Most interesting with regard to a prediction of the colour of the corresponding indigo dyes is a comparison of the longest wavelength absorption maxima of **6** and **7** with those of **2a-c**²⁴ (Table 2). Independent of the solvent used, **6** and **7** absorb approximately 25 and 50 nm, respectively, at longer wavelengths than **2c**. While the longest wavelength absorption maxima of **6** reach those of **2a** and **2b**, those of **7** surpass all. It could therefore well be, that the unknown indigo dye **8a**, containing the new chromophore **7**, albeit in six-membered rings, absorbs at longer wavelengths than the classical indigo dyes **1a-c**. The same could be true for **8b**.

Table 2. Longest wavelength absorption maxima of **2a-c**, **6** and **7** in different solvents.

	λ_{\max} [nm] (log ϵ)				
	2a	2b	2c	6	7
Cyclohexane	473 (4.05)	476 (4.17)	450 (4.13)	475 (3.76)	503 (3.74)
Ethanol	480 (3.99)	478 (4.07)	453 (4.09)	475 (3.76)	503 (3.72)
Benzene	483 (4.05)	480 (4.12)	454 (4.10)	478 (3.77)	507 (3.74)
Chloroform	487 (4.00)	482 (4.04)	458 (4.07)	481 (3.74)	510 (3.72)



8a X = S
8b X = Se

The PE spectrum of **7** is depicted in Figure 2; the relevant ionization potentials are summarized in Table 3 together with the results of quantum chemical calculations. In the low-energy region (< 11 eV), five ionization bands are found of which only one (that centred at 9.36 eV) seems to be composed of two different ionizations. Based on PM3 and B3LYP calculations, assignments are readily made. Compared to **2c**,^{9,10} the first IP is raised by 0.23 eV while the second is lowered by 0.33 eV.




Figure 2. PE spectrum of **7**

Table 3. Vertical ionization potentials IP_v [eV] and orbital energies ϵ [eV] of **7**.

IP_v exp.	PM3 - ϵ	B3LYP/6-31+G* - ϵ	$IP_v^{[a]}$	Assignment
	1.74	3.21		π_6 (LUMO)
7.83	8.64	5.96	7.53	π_5 (HOMO)
8.54	9.32	6.57	8.14	π_4
9.36	10.90	7.51	9.08	π_3
9.5 sh	10.95	7.66	9.23	$n^+(O)$
9.97	11.54	8.00	9.57	$n^-(O)$
11.0	12.26	9.10	10.67	$n^-(S)$
	12.37	9.36	10.93	$n^+(S)$

[a] Calculation of first vertical IP : energy difference of molecule ($E = -2131.51089$ au) and radical cation with identical geometry ($E = -2131.23399$ au). Higher IP s: $IP_i = -\epsilon_i + 1.57$ eV.

In accord with their low excitation energies and long-wavelength absorptions, indigoid compounds are characterized by narrow HOMO-LUMO splittings.¹⁰ The HOMO-LUMO energy gap of **7** is calculated as 6.90 eV (PM3) or 2.75 eV (B3LYP) which is lower than that obtained for **2c** (7.40 or 3.05 eV).¹⁰ The fact that **7** has a lower excitation energy than **2c** is reflected by the smaller HOMO-LUMO gap and can first of all be traced back to a larger stabilization of the LUMO than that of the HOMO.

EXPERIMENTAL

The PE spectrum was recorded on a UPG200 spectrometer of Leybold-Heraeus equipped with a He(I) radiation source (21.21 eV). Semi-empirical PM3¹¹ calculations were performed with the MOPAC93¹² program package, B3LYP¹³ calculations with the program GAUSSIAN 94.¹⁴ IR spectra were recorded on a Perkin-Elmer 225 spectrophotometer. UV/VIS spectra were obtained with a Cary Instruments spectrometer model 14. ¹H and ¹³C NMR spectra were measured with a Varian VXR 200 or a Bruker AMX 300 spectrometer. For standards other than TMS the following chemical shifts were used: $\delta_H(\text{CHCl}_3) = 7.24$ ppm, $\delta_C(\text{CDCl}_3) = 77.00$ ppm. Mass spectra were determined with a Varian MAT 311 A or 701 instrument operated at 70 eV. R_f values are quoted for Macherey & Nagel Polygram SIL G/UV254 plates. Melting points were observed on a Reichert microhot-stage and are not corrected.

3,3-Dimethyl-[1,2]dithiolan-4-one (4): A solution of sodium disulfide, prepared by heating sodium sulfide nonahydrate (48.0 g, 0.20 mol) and sulfur (6.40 g, 0.20 g atom) in ethanol (300 ml) for 1.5 h to reflux, was cooled to 30°C until a solution of **4** (48.8 g, 0.20 mol) in ethanol (150 ml) was added at such a rate that the internal temperature did not exceed 45°C (30 min). After the addition was complete, the mixture was heated for 1 h to reflux, cooled, poured into water (1.0 l) and extracted with dichloromethane (2 x 150 ml). The combined extracts were washed with water (2 x 150 ml), dried (MgSO₄) and concentrated on a rotary evaporator (bath temperature 30°C/15 torr). The residual brown oil (30 g) was fractionated in vacuo to yield 16.4 g (55%) **4** as an unpleasant smelling yellow liquid, bp 82–83°C/15 torr (lit^a 90–102°C/25 torr, lit^b 70°C/11 torr). ¹H NMR (300 MHz, CDCl₃, TMS int): 1.53 (s, 6H), 3.58 (s, 2H). ¹³C NMR (50 MHz, CDCl₃, CDCl₃ int): 23.70 (C_{prim}), 41.58 (C_{sek}), 55.70, 210.22 (C_{quar}). MS m/z 148 (100, M⁺). Anal. calcd for C₅H₈OS₂: C, 40.51; H, 5.44. Found: C, 41.63; H, 5.74.

Trans-4,4,4',4'-tetramethyl-[2,2']bithiolanylidene-3,3'-dione (2c), trans-5-(4,4-dimethyl-3-oxo-thiolan-2-ylidene)-3,3-dimethyl-[1,2]dithiolan-4-one (6) and trans-5,5,5',5'-tetramethyl-[3,3']bi[1,2']dithiolanylidene-4,4'-dione (7): To a solution of potassium hexacyanoferrate (III) (39.5 g, 120 mmol) and potassium hydroxide (15.6 g, 86%, 240 mmol) in water (400 ml) was added all at once a 1:1-mixture of **4** (2.22 g, 15 mmol) and **5** (1.95 g, 15 mmol) in ethanol (40 ml) causing a slightly exothermic effect and an immediate separation of an oily solid. The mixture was heated for 10 min on a water bath and then filtered. The residue was dissolved in dichloromethane (200 ml), and the solution was washed with water (2 x 100 ml) and dried (MgSO₄). The solvent was distilled off on a rotary evaporator (bath temperature 30°C/15 torr) and the remaining material (2.1 g) was chromatographed on silica gel (0.05–0.20 mm) in benzene [column 80 x 5 cm; R_f = 0.72 (**7**), 0.50 (**6**), 0.26 (**2c**)] yielding 590 mg (15%) **2c**, 430 mg (10%) **6** and 240 mg (5%) **7** as orange, light-red and deep-red solids, respectively. Analytically pure samples of **2c**, **6** and **7** were obtained by crystallization from ethanol. Oxidation of pure **4** gave only **7** (16%).

2c: Fp 232–234°C (lit⁴ 233–234°C). The ¹H NMR data were identical with literature data.⁴ The ¹³C NMR data have not yet been reported and were as follows: ¹³C NMR (50 MHz, CDCl₃, CDCl₃ int): δ = 24.33 (C_{prim}), 41.62 (C_{sek}), 45.97, 131.01, 207.36 (C_{quar}).

6: Fp 144°C (subl. 110°C). IR (KBr): C=O 1690, 1673 cm⁻¹. UV/VIS λ_{max} [mμ] (log ε): C₆H₁₂: 475, 376 sh, 358 (3.76, 3.41, 3.53), C₂H₅OH: 475, 375 sh, 357 (3.76, 3.43, 3.51), C₆H₆: 478, 377 sh, 359 (3.77, 3.44, 3.53), CHCl₃: 481, 377 sh, 359 (3.74, 3.42, 3.51), KBr: 495. ¹H NMR (300 MHz, CDCl₃, CHCl₃ int): δ = 1.28 (s, 6H), 1.57 (s, 6H), 3.10 (s, 2H). ¹³C NMR (50 MHz, CDCl₃, CDCl₃ int): δ = 24.06, 24.30 (C_{prim}), 42.35 (C_{sek}), 46.21, 56.77, 128.66, 137.25, 201.81, 207.40 (C_{quar}). MS m/z 274 (81, M⁺), 172 (100), 88 (25). Anal. calcd for C₁₁H₁₄O₃S₃: C, 48.14; H, 5.14; S, 35.05. Found: C, 47.93; H, 5.06; S, 35.30.

7: Fp 191°C (subl. 130°C). IR (KBr): C=O 1667 cm⁻¹. UV/VIS λ_{max} [mμ] (log ε): C₆H₁₂: 503, 386 sh, 367, 346, 295 (3.74, 3.02, 3.13, 3.20, 3.53), C₂H₅OH: 503, 387 sh, 364, 347, 295 (3.72, 3.02, 3.13, 3.19, 3.53), C₆H₆: 507, 386 sh, 368, 348, 296 (3.74, 3.00, 3.13, 3.20, 3.54), CHCl₃: 510, 386 sh, 368, 347, 296 (3.72, 2.98, 3.11, 3.20, 3.52), KBr: 553. ¹H NMR (300 MHz, CDCl₃, CHCl₃ int): δ = 1.60 (s). ¹³C NMR (75 MHz, CDCl₃, CDCl₃ int): δ = 24.06 (C_{prim}), 57.31, 134.76, 201.88 (C_{quar}). MS m/z 292 (100, M⁺), 190 (60), 106 (46). Anal. calcd for C₁₀H₁₂S₄: C, 41.07; H, 4.14; S, 43.86. Found: C, 41.03; H, 4.11; S, 44.10.

Crystal structure analyses: **6:** C₁₀H₁₄O₂S₃, *Mr* = 274.40, crystal size: 0.40 × 0.40 × 0.30 mm³, monoclinic, space group *P*2₁/*n*, *a* = 8.4162(1), *b* = 8.1001(9), *c* = 9.209(1) Å, β = 100.953(7)°, *V* = 616.3(1) Å³, *Z* = 2, ρ_{calcd} = 1.479 Mg m⁻³, *F*(000) = 288, λ = 0.71073 Å, *T* = 133 K, μ(MoKα) = 0.583 mm⁻¹. Total number of reflections measured 13765, unique 1081 (*R*_{int} = 0.0288). Data/restraints/parameters: 1081/69/85, data collection range: 3.01 ≤ θ ≤ 25.03°. Final *R* indices: *R*1 = Σ|*F*_o - *F*_c|| / Σ|*F*_o| = 0.0602, *wR*2 = [Σ*w*(*F*_o² - *F*_c²)² / Σ*wF*_o⁴]^{1/2} = 0.1437 on data with *I* > 2σ(*I*) and *R*1 = 0.0605, *wR*2 = 0.1438 on all data; goodness of fit *S* = [Σ*w*(*F*_o² - *F*_c²)² / Σ(*n*-*p*)]^{1/2} = 1.372; extinction coefficient 0.015(4); largest difference peak and hole: 0.453 and -0.580 e·Å⁻³. **7:** C₁₉H₁₂O₂S₄, *Mr* = 292.44, crystal size: 0.70 × 0.50 × 0.20 mm³, monoclinic, space group *P*2₁/*n*, *a* = 8.265(2), *b* = 8.228(2), *c* = 9.178(2) Å, β = 101.14(3)°, *V* = 612.4(2) Å³, *Z* = 2, ρ_{calcd} = 1.586 Mg m⁻³, *F*(000) = 304, λ = 0.71073 Å, *T* = 153 K, μ(MoKα) = 1.495 mm⁻¹. Total number of reflections measured 10502, unique 1248 (*R*_{int} = 0.0366). Data/restraints/parameters: 1248/0/76, data collection range: 3.04 ≤ θ ≤ 26.37°. Final *R* indices: *R*1 = 0.0221, *wR*2 = 0.0578 on data with *I* > 2σ(*I*) and *R*1 = 0.0245, *wR*2 = 0.0590 on all data; goodness of fit *S* = 1.082; extinction coefficient 0.060(4); largest difference peak and hole: 0.398 and -0.230 e·Å⁻³. The crystals were mounted on a glass fiber in a rapidly cooled perfluoropolyether.¹⁵ Diffraction data were collected on a Stoe-Siemens-Huber four-circle-diffractometer coupled to a Siemens CCD area-detector at 133(2) K, with graphite-monochromated MoKα radiation (λ = 0.71073 Å), performing φ- and ω-scans. The

structures were solved by direct methods using the program SHELXS-97¹⁶ and refined against F^2 on all data by full-matrix least squares with SHELXL-97.¹⁷ All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in the model at geometrically calculated positions and refined using a riding model. The disorder in **6** was modelled with the help of similarity restraints for 1-2 and 1-3 distances and displacement parameters as well as rigid bond restraints for anisotropic displacement parameters. Crystallographic data for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC 133014 and CCDC 133015, respectively. Copies of the data can be obtained free of charge on application to the CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: Code + (44) 1223 336-033; e-mail: deposit@ccdc.cam.ac.uk).

REFERENCES AND NOTES

Dedicated to Professor Wolfgang Lüttke on the occasion of his 80th birthday

- 1 (a) Klessinger, M.; Lüttke, W. *Tetrahedron* **1963**, *19*, Suppl. 2, 315-355. (b) Klessinger, M. *Tetrahedron* **1966**, *22*, 3355-3365.
- 2 **2a**: (a) Wille, E.; Lüttke, W. *Angew. Chem.* **1971**, *83*, 853-854; *Angew. Chem. Int. Ed. Engl.* **1971**, *10*, 803-804. (b) Wille, E.; Lüttke, W. *Liebigs Ann. Chem.* **1980**, 2039-2054.
- 3 **2b**: Fitjer, L.; Lüttke, W. *Chem. Ber.* **1972**, *105*, 919-928.
- 4 **2c**: (a) Lüttke, W.; Hermann, H.; Klessinger, M. *Angew. Chem.* **1966**, *78*, 638-639; *Angew. Chem. Int. Ed. Engl.* **1966**, *5*, 598-599. (b) Hermann, H.; Lüttke, W. *Chem. Ber.* **1968**, *101*, 1708-1714. (c) Hermann, H.; Lüttke, W. *Chem. Ber.* **1968**, *101*, 1715-1728.
- 5 Open-chain chromophores derived from **1c** are nonplanar and absorb at distinctly shorter wavelengths than **2c**: Hermann, H.; Lüttke, W. *Chem. Ber.* **1971**, *104*, 492-512.
- 6 Fitjer, L.; Lüttke, W. *Chem. Ber.* **1972**, *105*, 907-918.
- 7 A formation of **5** has formerly been observed during treatment of **4** with (a) sodium sulfide (Luhmann, U.; Wentz, F. G.; Lüttke, W.; Süsse, P. *Chem. Ber.* **1976**, *110*, 1421-1431) and (b) sodium hydrogen-sulfide (Föhlich, B.; Gottstein, W. *Liebigs Ann. Chem.* **1979**, 1768-1784).
- 8 Wagner, R. B.; Moore, J. A. *J. Am. Chem. Soc.* **1950**, *72*, 974-977.
- 9 Bauer, H.; Kowski, K.; Kuhn, H.; Lüttke, W.; Rademacher, P. *J. Mol. Struct.* **1998**, *445*, 277-286.
- 10 Rademacher, P.; Kowski, K.; Hermann, H.; Lüttke, W. *Eur. J. Org. Chem.* **1999**, submitted.
- 11 Stewart, J. J. P. *J. Comput. Chem.* **1989**, *10*, 209-220, 221-264.
- 12 Stewart, J. J. P. *MOPAC93.00 Manual*, Fujitsu Limited, Tokyo, Japan, **1993**.
- 13 Becke, A. D. *J. Comput. Chem.* **1999**, *20*, 63-69; and references given therein.
- 14 Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. P.; Head-Gordon, M.; Gonzales, C.; Pople, J. A. *Gaussian 94, Revision E.2*, Gaussian, Inc., Pittsburgh, PA, **1995**.
- 15 Kottke, J.; Stalke, D. *J. Appl. Crystallogr.* **1993**, *26*, 615-619.
- 16 Sheldrick, G. M. *Acta Crystallogr., Sect. A*, **1990**, *46*, 467-473.
- 17 Sheldrick, G. M. SHELX 97, Universität Göttingen, 1997.